

Important Medication Information for Allergy Testing.

Medications can interfere with allergy skin testing. The following list of medications need to be stopped the specified number of days prior to your visit.

DO NOT STOP TAKING YOUR ASTHMA MEDICATIONS

STOP DAYS		STOP DAYS		STOP DAYS	
2	Actifed	5	Clemastine (RX)	2	Nizatidine
2	Advil-PM	3	Compazine	2	Nyquil
5	Alavert	2	Contac	2	Olopatadine (nasal)
5	Allegra (180 mg)	5	CyClobenzaprine	2	Pataday
2	Allegra (30 mg, 60 mg)	5	CyProheptadine	2	Patanase
2	Allegra-D (12-Hour)	6	Desloratadine	2	Pepcid
5	Allegra-D (24-hour)	5	Dexchlorpheniramine	5	Periactin
3	Antivert	3	Dimenhydrinate	3	Phenergan
5	Arbinoxa	2	Dimetane	3	Prochlorperazine
2	Astelin	2	Dimetapp	3	Promethazine
5	Atarax	2	Diphenhydramine	2	Sominex
2	Axid	3	Dramamine	2	Tagamet
2	Azelastine	2	Dymista	5	Tavist (non-drowsy)
2	Benadryl	2	Excedrin-PM	2	Tavist (OTC)
2	Bromfed	2	Famotidine	5	Tavist (RX)
2	Brompheniramine	5	Fexofenadine (180 mg)	3	Tigan
5	Carbinoxamine	2	Fexofenadine (low-dose)	3	Trimethobenzamide
5	Cetirizine	5	Flexeril	2	Tripolidine
2	Chlorpheniramine	2	Histex	2	Tums Ultra
2	Chlor-Trimeton	5	Hydroxyzine	2	Tylenol-PM
2	Cimetidine	5	Karbinal	5	Vistaril
6	Clarinet	2	Kronofed	5	Xyzal
6	Clarinet-D	5	Levocetirizine	2	Z-Quil
5	Claritin	5	Loratadine products	5	Zyrtec
2	Clemastine (OTC)	3	Meclizine	5	Zyrtec-D

● **DECONGESTANTS:** Stop taking only when combined with antihistamines.

● **ANTACIDS:** You can continue taking: *Prevacid, *Prilosec, *Nexium and other "PPI" drugs (*lansoprazole, *omeprazole, etc).